Finsleroid-regular space: curvature tensor, continuation of gravitational Schwarzschild metric

G.S. Asanov

Division of Theoretical Physics, Moscow State University
119992 Moscow, Russia
(*e-mail: asanov@newmail.ru*)

arXiv:0712.0440v1 [math-ph] 4 Dec 2007

Abstract

The method of simple straightforward calculation of the curvature tensor of the Finsleroid–regular space is indicated. The Schwarzschild metric which underlines the gravitational field produced by static spherical-symmetric body is shown to be uniquely extended to the Finslerian domain upon a consistent treatment of the pseudo-Finsleroid axis vector field b_i to be the field of the time variable.

Keywords: Finsler metrics, gravitational equations, curvature tensors.

1. Description of new conclusions

The regular Finsleroid-Finsler spaces [1] of the positive-definite type $\mathcal{FR}^{PD}_{g;c}$ and of the relativistic case $\mathcal{FR}^{SR}_{g;c}$ are constructed with the help of the set $\{g(x), b_i(x), a_{ij}(x), y\}$, where $g(x)$ is a scalar which plays the role of the Finsleroid charge, $b_i(x)$ is an involved vector field, and $a_{ij}(x)$ is a metric tensor (which is positive-definite in the space $\mathcal{FR}^{PD}_{g;c}$ and indefinite in the space $\mathcal{FR}^{SR}_{g;c}$). The norm

$$||b|| = c \equiv \sqrt{a_{ij}(x)b^i(x)b^j(x)} \tag{1.1}$$

of the input 1-form $b = b_i y^i$ is taken to be an arbitrary positive scalar subject to the restrictions: $0 < c(x) < 1$ in the positive-definite space $\mathcal{FR}^{PD}_{g;c}$, and $c(x) > 1$ in the indefinite space $\mathcal{FR}^{SR}_{g;c}$. In the latter space, we assume the 1-form b to be time-like.

If we specify the Riemannian metric tensor according to

$$a_{ij} = \frac{1}{c^2} b_i b_j + m u_{ij} \tag{1.2}$$

(with $b^i u_{ij} = 0$ and $b^i = a^{ij} b_j$; u_{ij} is the $(N-1)$-dimensional part of Euclidean metric tensor, in compliance with (A.1)-(A.7)) and assume the scalars c, m to be of the dependence $c = c(r)$, $m = m(r)$ on the radius variable $r = \sqrt{u_{ij}x^i x^j}$ (assuming also $c > 0$ and $m \neq 0$), we can readily conclude that the Riemannian covariant derivative

$$\nabla_i b_j = \frac{\partial b_j}{\partial x^i} - b_n a^n{}_{ij} \tag{1.3}$$

($a^n{}_{ij}$ stand for the Christoffel symbols constructed from the tensor $a_{ij}(x)$) fulfills the equality

$$\nabla_i b_j = \frac{1}{c}(c_i b_j + c_j b_i) \tag{1.4}$$

(the respective calculations are presented in Appendix A); $c_i = \partial c/\partial x^i$. The right-hand part of (1.4) does not involve the function m. It follows that

$$y^i y^j \nabla_i b_j = \frac{2}{c} b(yc), \tag{1.5}$$

where $(yc) = y^h c_h$. We have

$$b^i c_i = 0. \tag{1.6}$$

Under the condition (1.2) *the Finsleroid-regular spaces cannot be of neither Berwald case nor the Landsberg type.* Obviously, the Landsberg characteristic condition $\nabla_i b_j = k r_{ij}$ (derived and used in [2-5], and applied to cosmological consideration in [6]) is not realized when (1.4) takes place. The Berwald case would imply $\nabla_i b_j = 0$ (see [1]), that is $c_i = 0$.

At the same time, the Schwarzschild metric (which underlines the gravitational field produced by static spherical-symmetric body) belongs to the form (1.2). Therefore, the pseudo-Finsleroid regular space $\mathcal{FR}^{SR}_{g;c}$ taken together with the representations (1.2)-(1.4) provides us with a straightforward continuation of the Riemannian Schwarzschild-metric framework into the pseudo-Finslerian domain with respect to the pseudo-Finsleroid charge $g(x)$ (which thereby plays the role of the characteristic parameter of extension). The pseudo-Finsleroid axis vector field $b_i(x)$ plays the role of the distribution of the time variable.

Key objects involved are explicitly calculated below in Appendix A upon using the metric tensor (1.2). Namely, calculating the associated Christoffel symbols ${a^k}_{ij}$ leads to (A.31), thereafter the Riemannian curvature tensor ${a_n}^i{}_{km}$ (see the definition (A.32)) is obtained according to (A.39), and the entailed Ricci tensor ${a_n}^i{}_{im}$ is given by (A.40)-(A.43). With these representations, we examine the particular case:

$$c = \frac{1 + \dfrac{\xi}{4r}}{1 - \dfrac{\xi}{4r}} \tag{1.7}$$

(see (A.47)) and

$$m = -\left(1 + \frac{\xi}{4r}\right)^4 \tag{1.8}$$

(see (A.49)), obtaining

$${a_n}^i{}_{im} = 0 \quad \text{when} \quad N = 4. \tag{1.9}$$

The respective curvature tensor ${a_n}^i{}_{km}$ is found to read (A.54). The vanishing (1.9) demonstrates that the metric tensor (1.2) with the choice (1.7) and (1.8) fulfills the gravitational field equations in vacuum. Raising forth the identification

$$\xi = r_g, \tag{1.10}$$

where r_g is the gravitational radius of the massive static and spherical-symmetric body, we may conclude that in the dimension $N = 4$ the metric tensor (1.2) with the choice (1.7) and (1.8) is just the Schwarzschild metric tensor (cf. [8,9]).

It should be noted that $c < 1$ of the positive-definite space transforms to $c > 1$ in the relativistic counterpart with the timelike vector b_i. The reason is that the Riemannian metric

$$S^2 = b^2 + q^2$$

of the positive-definite space must be replaced by

$$S^2 = b^2 - q^2,$$

when treating the relativistic case.

In Appendix B, using the spray coefficients found in the previous work [1], we evaluate the associated Finslerian curvature tensor ${R^i}_k$ under the only conditions that $\nabla_n b_m - \nabla_m b_n = 0$ and $g = const$, simultaneously keeping the general case $c = c(x)$ of the norm value $||b||$. The resultant representation is sufficiently simple and transparent and, therefore, opens up the operative possibility to evaluate the tensor

$$\rho_{ij} := \frac{1}{2}({R_i}^m{}_{mj} + R^m{}_{ijm}) - \frac{1}{2} g_{ij} R^{mn}{}_{nm} \tag{1.11}$$

and the current

$$\rho^i := {\rho^i}_j y^j \tag{1.12}$$

(see [2,6]) which can be used to represent the energy-momentum distribution in the respective pseudo-Finsleroid-regular space-time. Various relativistic and other applications are possible.

Appendix A: Spherical-symmetric Riemannian background

Let us start with the (pseudo)Euclidean metric tensor e_{ij}, which inverse will be denoted by e^{ij}, so that $e_{ij}e^{jn} = \delta_i{}^n$, and introduce the decomposition

$$e_{ij} = e_i e_j + \epsilon u_{ij}, \qquad \epsilon = \pm 1, \tag{A.1}$$

where e_i is a unit vector and

$$\mathrm{rank}(u_{ij}) = N - 1, \qquad y^i y^j u_{ij} \geq 0. \tag{A.2}$$

The value $\epsilon = 1$ corresponds to the positive-definite case; in the relativistic case we must take $\epsilon = -1$. If we construct the contravariant vector

$$e^i = e^{ij} e_j, \tag{A.3}$$

we obtain

$$e_i e^i = 1, \qquad e^i u_{ij} = 0. \tag{A.4}$$

We get also

$$u_i{}^j = u^{jm} u_{im}, \qquad u^{ij} = e^{im} e^{jn} u_{mn}, \tag{A.5}$$

and

$$u^{ij} u_{jn} = \delta^i{}_n - e^i e_n, \qquad u_i{}^j = \delta_i{}^j - e_i e^j. \tag{A.6}$$

Let us set forth the identification

$$b_i = e_i, \tag{A.7}$$

introduce two scalars $c > 0$ and $m \neq 0$, and specify the Riemannian metric tensor as follows:

$$a_{ij} := \frac{1}{c^2} b_i b_j + m u_{ij}, \tag{A.8}$$

which inverse reads

$$a^{ij} = \frac{1}{c^2} b^i b^j + \frac{1}{m} u^{ij}, \tag{A.9}$$

so that $a_{ij} a^{jn} = \delta_i{}^n$.

We use the contravariant vector

$$b^i = a^{ij} b_j, \tag{A.10}$$

which entails

$$b^i u_{ij} = 0 \tag{A.11}$$

and

$$b^i b_i = c^2, \tag{A.12}$$

together with

$$b^i = c^2 e^i. \tag{A.13}$$

We introduce the radius variable

$$r = \sqrt{u_{ij} x^i x^j}, \tag{A.14}$$

obtaining

$$n_i \ := \frac{1}{r} u_{ij} x^j = \frac{\partial r}{\partial x^i}, \tag{A.15}$$

$$\frac{\partial n_i}{\partial x^j} = \frac{1}{r} \left(u_{ij} - n_i n_j \right), \tag{A.16}$$

$$n^i = u^{ij} n_j = \epsilon e^{ij} n_j, \tag{A.17}$$

and also

$$n^i n_i = 1 \tag{A.18}$$

together with

$$n_i b^i = 0. \tag{A.19}$$

We shall specify the scalars c, m to be of the dependence

$$c = c(r), \qquad m = m(r), \tag{A.20}$$

assuming $c > 0$ and $m \neq 0$. We shall also use the notation

$$c_i = \frac{\partial c}{\partial x^i} \equiv c' n_i \tag{A.21}$$

and

$$c^i = a^{ij} c_j, \tag{A.22}$$

so that

$$c^i = \frac{1}{m} c' n^i \tag{A.23}$$

and

$$b^i c_i = 0. \tag{A.24}$$

Under these conditions, we obtain

$$\frac{\partial a_{ij}}{\partial x^n} = n_n \left(-\frac{2}{c^3} c' b_i b_j + m' u_{ij} \right), \tag{A.25}$$

from which it just follows that

$$b_n a^n{}_{ij} = \frac{1}{2} b^n \left[n_i \left(-\frac{2c'}{c^3} b_n b_j + m' u_{nj} \right) + n_j \left(-\frac{2}{c^3} c' b_i b_n + m' u_{in} \right) + n_n \left(\frac{2}{c^3} c' b_i b_j - m' u_{ij} \right) \right],$$

or

$$b_n a^n{}_{ij} = -\frac{1}{c} c' (n_i b_j + n_j b_i) = -\frac{1}{c} (c_i b_j + c_j b_i).$$

Thus for the Riemannian covariant derivative

$$\nabla_i b_j \ = \frac{\partial b_j}{\partial x^i} - b_n a^n{}_{ij} \tag{A.26}$$

we obtain the simple expression

$$\nabla_i b_j = \frac{1}{c} (c_i b_j + c_j b_i), \tag{A.27}$$

the right-hand part of which does not involve the function m. It follows that

$$y^i y^j \nabla_i b_j = \frac{2}{c} b(yc). \tag{A.28}$$

We use the notation

$$(yc) = y^h c_h. \tag{A.29}$$

With (A.25), we also obtain

$$c_n a^n{}_{ij} = \frac{1}{2} c^n \left[n_i \left(-\frac{2}{c^3} c' b_n b_j + m' u_{nj} \right) + n_j \left(-\frac{2}{c^3} c' b_i b_n + m' u_{in} \right) + n_n \left(\frac{2}{c^3} c' b_i b_j - m' u_{ij} \right) \right]$$

and derive

$$c_n a^n{}_{ij} = \frac{1}{2} \frac{1}{m} c' \left[2m' n_i n_j + \frac{2}{c^3} c' b_i b_j - m' u_{ij} \right],$$

so that

$$\nabla_i c_j = c'' n_i n_j + c' \frac{1}{r} (u_{ij} - n_i n_j) - \frac{1}{2} \frac{1}{m} c' \left[2m' n_i n_j + \frac{2}{c^3} c' b_i b_j - m' u_{ij} \right]. \tag{A.30}$$

Calculating the Christoffel symbols $a^k{}_{ij}$ on the basis of (A.8) and (A.9) yields straightforwardly that

$$a^k{}_{ij} = \frac{1}{2} a^{kn} \left[n_i \left(-\frac{2}{c^3} c' b_n b_j + m' u_{nj} \right) + n_j \left(-\frac{2}{c^3} c' b_n b_i + m' u_{ni} \right) + n_n \left(\frac{2}{c^3} c' b_i b_j - m' u_{ij} \right) \right],$$

or

$$a^k{}_{ij} = -\frac{1}{c^3} c' b^k (n_i b_j + n_j b_i) + \frac{1}{2} \frac{1}{m} \left[m' n_i u_j{}^k + m' n_j u_i{}^k + n^k \left(\frac{2}{c^3} c' b_i b_j - m' u_{ij} \right) \right]. \tag{A.31}$$

We are aimed now to obtain the Riemannian curvature tensor

$$a_n{}^i{}_{km} = \frac{\partial a^i{}_{nm}}{\partial x^k} - \frac{\partial a^i{}_{nk}}{\partial x^m} + a^u{}_{nm} a^i{}_{uk} - a^u{}_{nk} a^i{}_{um}. \tag{A.32}$$

To this end we write (A.31) in the form

$$a^i{}_{nm} = -\frac{1}{c} c' e^i (n_n b_m + n_m b_n) + \frac{1}{2} \frac{1}{m} \left[m' n_n u_m{}^i + m' n_m u_n{}^i + n^i \left(\frac{2}{c^3} c' b_n b_m - m' u_{nm} \right) \right] \tag{A.33}$$

and get

$$\frac{\partial a^i{}_{nm}}{\partial x^k} - \frac{\partial a^i{}_{nk}}{\partial x^m} = -\left(\frac{1}{c} c'' - \frac{1}{c^2} c' c' \right) e^i n_n (n_k b_m - n_m b_k)$$

$$-\frac{1}{c} c' \frac{1}{r} \Big[(u_{nk} - n_n n_k)\, b_m - (u_{nm} - n_n n_m)\, b_k \Big] e^i$$

$$-\frac{1}{2}\frac{1}{m^2}m'n_k\left[m'n_n u_m{}^i - n^i\left(-\frac{2}{c^3}c'b_n b_m + m'u_{nm}\right)\right]$$

$$+\frac{1}{2}\frac{1}{m^2}m'n_m\left[m'n_n u_k{}^i - n^i\left(-\frac{2}{c^3}c'b_n b_k + m'u_{nk}\right)\right]$$

$$+\frac{1}{2}\frac{1}{m}\left[m''n_n n_k + m'\frac{1}{r}\left(u_{nk} - n_n n_k\right)\right]u_m{}^i$$

$$-\frac{1}{2}\frac{1}{m}\left[m''n_n n_m + m'\frac{1}{r}\left(u_{nm} - n_n n_m\right)\right]u_k{}^i$$

$$+\frac{1}{2}\frac{1}{m}\frac{1}{r}(u^i{}_k - n^i n_k)\left(\frac{2}{c^3}c'b_n b_m - m'u_{nm}\right)$$

$$-\frac{1}{2}\frac{1}{m}\frac{1}{r}(u^i{}_m - n^i n_m)\left(\frac{2}{c^3}c'b_n b_k - m'u_{nk}\right)$$

$$+\frac{1}{m}n^i\left(\frac{1}{c^3}c'' - \frac{3}{c^4}c'c'\right)b_n(n_k b_m - n_m b_k) - \frac{1}{2}\frac{1}{m}m''(n_k u_{nm} - n_m u_{nk})n^i,$$

or

$$\frac{\partial a^i{}_{nm}}{\partial x^k} - \frac{\partial a^i{}_{nk}}{\partial x^m} = -\frac{m'}{m}\frac{1}{r}\left(u_{mn}u_k{}^i - u_{kn}u_m{}^i\right)$$

$$-\frac{1}{c^2}\left(\frac{1}{c}c'' - \frac{1}{c^2}c'c' - \frac{1}{r}\frac{1}{c}c'\right)n_n(n_k b_m - n_m b_k)b^i$$

$$-\frac{1}{c^3}c'\frac{1}{r}(u_{nk}b_m - u_{nm}b_k)b^i$$

$$+\frac{1}{2}\frac{1}{m}\frac{m'}{m}\left[n_k\left(-\frac{2}{c^3}c'b_n b_m + m'u_{nm}\right) - n_m\left(-\frac{2}{c^3}c'b_n b_k + m'u_{nk}\right)\right]n^i$$

$$+\frac{1}{2}\frac{1}{m}\left[m'' - \frac{1}{r}m' - \frac{(m')^2}{m}\right]n_n\left(n_k u_m{}^i - n_m u_k{}^i\right)$$

$$+\frac{1}{m}\frac{1}{r}\frac{1}{c^3}c'b_n\left(b_m u^i{}_k - b_k u^i{}_m\right)$$

$$+\frac{1}{m}\frac{1}{c^2}\left(\frac{1}{c}c'' - \frac{3}{c^2}c'c' - \frac{1}{r}\frac{1}{c}c'\right)b_n(n_k b_m - n_m b_k)n^i$$

$$-\frac{1}{2}\frac{1}{m}\left(m'' - \frac{1}{r}m'\right)(n_k u_{nm} - n_m u_{nk})n^i.$$

In this way we get

$$\frac{\partial a^i{}_{nm}}{\partial x^k} - \frac{\partial a^i{}_{nk}}{\partial x^m} = -\frac{m'}{m}\frac{1}{r}\left(u_{mn}u_k{}^i - u_{kn}u_m{}^i\right)$$

$$-\frac{1}{c^2}\left(\frac{1}{c}c'' - \frac{1}{c^2}c'c' - \frac{1}{r}\frac{1}{c}c'\right)\left[n_n(n_kb_m - n_mb_k)b^i - \frac{1}{m}b_n(n_kb_m - n_mb_k)n^i\right]$$

$$-\frac{1}{c^3}c'\frac{1}{r}(u_{nk}b_m - u_{nm}b_k)b^i - \frac{1}{m}\frac{m'}{m}\frac{1}{c^3}c'b_n(n_kb_m - n_mb_k)n^i$$

$$-\frac{1}{2}\frac{1}{m}\left[m'' - \frac{1}{r}m' - \frac{(m')^2}{m}\right]\left(n_n(n_mu_k{}^i - n_ku_m{}^i) - (n_mu_{nk} - n_ku_{nm})n^i\right)$$

$$+\frac{1}{m}\frac{1}{r}\frac{1}{c^3}c'b_n\left(b_mu^i{}_k - b_ku^i{}_m\right) - \frac{1}{m}\frac{1}{c^2}\frac{2}{c^2}c'c'b_n(n_kb_m - n_mb_k)n^i. \qquad \text{(A.34)}$$

Next, we obtain the representation

$$a^u{}_{nm}a^i{}_{uk} - a^u{}_{nk}a^i{}_{um} = -\frac{1}{c}c'(n_nb_m + n_mb_n)\left[-\frac{1}{c}c'n_ke^i + \frac{1}{m}\frac{1}{c^3}c'b_kn^i\right]$$

$$-\frac{1}{2}\frac{1}{c}c'\frac{1}{m}m'n_nn_mb_ke^i$$

$$+\frac{1}{2}\frac{m'}{m}n_mu_n{}^u\left[-\frac{c'}{c}e^i(n_ub_k + n_kb_u) + \frac{1}{2}\frac{1}{m}\left(m'n_uu_k{}^i + m'n_ku_u{}^i + n^i\left(\frac{2c'}{c^3}b_ub_k - m'u_{uk}\right)\right)\right]$$

$$+\frac{1}{2}\frac{1}{m}\left(\frac{2}{c^3}c'b_nb_m - m'u_{nm}\right)\left[-\frac{1}{c}c'e^ib_k + \frac{1}{2}\frac{1}{m}m'u_k{}^i\right] - [km]$$

$$= \frac{1}{c^2}c'c'n_n(b_mn_k - b_kn_m)e^i - \frac{1}{c^4}c'c'\frac{1}{m}b_n(n_mb_k - n_kb_m)n^i$$

$$-\frac{1}{2}\frac{1}{c}c'\frac{m'}{m}n_n(n_mb_k - n_kb_m)e^i$$

$$-\frac{1}{2}\frac{m'}{m}\frac{c'}{c}n_n(n_mb_k - n_kb_m)e^i + \frac{1}{4}\left(\frac{m'}{m}\right)^2\left(n_n(n_mu_k{}^i - n_ku_m{}^i) - (n_mu_{nk} - n_ku_{nm})n^i\right)$$

$$+\frac{1}{2}\frac{1}{m}\frac{c'}{c^3}\frac{m'}{m}b_n(b_mu_k{}^i - b_ku_m{}^i) - \frac{1}{2}\frac{m'}{m}u_{nm}\left[-\frac{c'}{c}e^ib_k + \frac{1}{2}\frac{m'}{m}u_k{}^i\right] + \frac{1}{2}\frac{m'}{m}u_{nk}\left[-\frac{c'}{c}e^ib_m + \frac{1}{2}\frac{m'}{m}u_m{}^i\right].$$

Eventually,

$$a^u{}_{nm}a^i{}_{uk} - a^u{}_{nk}a^i{}_{um} =$$

$$= \frac{1}{c^4}c'c'\left[n_n(b_m n_k - b_k n_m)b^i - b_n(n_m b_k - n_k b_m)\frac{1}{m}n^i\right]$$

$$-\frac{1}{c^3}c'\frac{m'}{m}n_n(n_m b_k - n_k b_m)b^i$$

$$+\frac{1}{4}\left(\frac{m'}{m}\right)^2\left(n_n(n_m u_k{}^i - n_k u_m{}^i) - (n_m u_{nk} - n_k u_{nm})n^i\right)$$

$$+\frac{1}{2}\frac{1}{m}\frac{c'}{c^3}\frac{m'}{m}b_n(b_m u_k{}^i - b_k u_m{}^i)$$

$$-\frac{1}{2}\frac{m'}{m}u_{nm}\left[-\frac{1}{c^3}c'b^ib_k + \frac{1}{2}\frac{m'}{m}u_k{}^i\right] + \frac{1}{2}\frac{m'}{m}u_{nk}\left[-\frac{1}{c^3}c'b^ib_m + \frac{1}{2}\frac{m'}{m}u_m{}^i\right]. \tag{A.35}$$

Adding (A.35) to (A.34) we find the curvature tensor (A.32):

$$a_n{}^i{}_{km} = -\frac{m'}{m}\left(\frac{1}{4}\frac{m'}{m} + \frac{1}{r}\right)\left(u_{mn}u_k{}^i - u_{kn}u_m{}^i\right)$$

$$-\frac{1}{c^2}\left(\frac{1}{c}c'' - \frac{2}{c^2}c'c' - \frac{1}{r}\frac{1}{c}c' - \frac{c'}{c}\frac{m'}{m}\right)\left[n_n(n_k b_m - n_m b_k)b^i - \frac{1}{m}b_n(n_k b_m - n_m b_k)n^i\right]$$

$$-\frac{1}{2}\frac{1}{m}\left[m'' - \frac{1}{r}m' - \frac{3}{2}\frac{(m')^2}{m}\right]\left(n_n(n_m u_k{}^i - n_k u_m{}^i) - (n_m u_{nk} - n_k u_{nm})n^i\right)$$

$$+\frac{1}{2}\frac{c'}{c^3}\left(\frac{m'}{m} + \frac{2}{r}\right)\left[\frac{1}{m}b_n(b_m u_k{}^i - b_k u_m{}^i) - (b_m u_{nk} - b_k u_{nm})b^i\right]. \tag{A.36}$$

Inserting here

$$m u_{mn} = a_{mn} - \frac{1}{c^2}b_m b_n \tag{A.37}$$

and

$$u_i{}^k = \delta_i{}^k - \frac{1}{c^2}b_i b^k \tag{A.38}$$

yields the result

$$a_n{}^i{}_{km} = -\frac{1}{m}\frac{m'}{m}\left(\frac{1}{4}\frac{m'}{m} + \frac{1}{r}\right)\left(a_{mn}\delta_k{}^i - a_{kn}\delta_m{}^i\right)$$

$$+\frac{1}{c^2}\frac{m'}{m}\left(\frac{1}{4}\frac{m'}{m}+\frac{1}{r}\right)\left(\frac{1}{m}(a_{mn}b_kb^i-a_{kn}b_mb^i)+b_mb_n\delta_k{}^i-b_kb_n\delta_m{}^i\right)$$

$$-\frac{1}{c^2}\left(\frac{1}{c}c''-\frac{2}{c^2}c'c'-\frac{1}{r}\frac{1}{c}c'-\frac{c'}{c}\frac{m'}{m}\right)\left[n_n(n_kb_m-n_mb_k)b^i-\frac{1}{m}b_n(n_kb_m-n_mb_k)n^i\right]$$

$$-\frac{1}{2}\frac{1}{m}\left[m''-\frac{1}{r}m'-\frac{3}{2}\frac{(m')^2}{m}\right]\left(n_n(n_mu_k{}^i-n_ku_m{}^i)-(n_mu_{nk}-n_ku_{nm})n^i\right)$$

$$+\frac{1}{2}\frac{c'}{c^3}\left(\frac{m'}{m}+\frac{2}{r}\right)\left[\frac{1}{m}b_n(b_mu_k{}^i-b_ku_m{}^i)-(b_mu_{nk}-b_ku_{nm})b^i\right]. \tag{A.39}$$

Contracting over two indices leads to the representation

$$a_n{}^i{}_{im}=-\frac{m'}{m}\left(\frac{1}{4}\frac{m'}{m}+\frac{1}{r}\right)(N-2)u_{nm}$$

$$+\frac{1}{c^2}\left(\frac{1}{c}c''-\frac{2}{c^2}c'c'-\frac{1}{r}\frac{1}{c}c'-\frac{c'}{c}\frac{m'}{m}\right)\left[c^2n_nn_m+\frac{1}{m}b_nb_m\right]$$

$$-\frac{1}{2}\frac{1}{m}\left(m''-\frac{1}{r}m'-\frac{3}{2}\frac{(m')^2}{m}\right)\left[u_{nm}+(N-3)n_nn_m\right]$$

$$+\frac{1}{2}\frac{c'}{c^3}\left(\frac{m'}{m}+\frac{2}{r}\right)\left[c^2u_{nm}+(N-1)\frac{1}{m}b_nb_m\right],$$

which can be written as

$$a_n{}^i{}_{im}=n_1u_{nm}+\frac{1}{c^2}\frac{1}{m}n_2b_mb_n+n_3n_mn_n \tag{A.40}$$

with

$$n_1=-(N-2)\frac{m'}{m}\left(\frac{1}{4}\frac{m'}{m}+\frac{1}{r}\right)-\frac{1}{2}\frac{1}{m}\left(m''-\frac{1}{r}m'-\frac{3}{2}\frac{(m')^2}{m}\right)+\frac{1}{2}\frac{c'}{c}\left(\frac{m'}{m}+\frac{2}{r}\right), \tag{A.41}$$

$$n_2=\left(\frac{1}{c}c''-\frac{2}{c^2}c'c'-\frac{1}{r}\frac{1}{c}c'-\frac{c'}{c}\frac{m'}{m}\right)+\frac{1}{2}\frac{c'}{c}\left(\frac{m'}{m}+\frac{2}{r}\right)(N-1), \tag{A.42}$$

and

$$n_3=\left(\frac{1}{c}c''-\frac{2}{c^2}c'c'-\frac{1}{r}\frac{1}{c}c'-\frac{c'}{c}\frac{m'}{m}\right)-\frac{1}{2}\frac{1}{m}\left(m''-\frac{1}{r}m'-\frac{3}{2}\frac{(m')^2}{m}\right)(N-3). \tag{A.43}$$

Let us make the substitution

$$c = y\left(\frac{\xi}{4r}\right), \qquad m = z\left(\frac{\xi}{4r}\right), \qquad \xi = const, \tag{A.44}$$

so that

$$c' = -\frac{\xi}{4r^2}y', \qquad m' = -\frac{\xi}{4r^2}z', \tag{A.45}$$

and

$$c'' = \frac{\xi}{2r^3}y' + \frac{\xi^2}{16r^4}y'', \qquad m'' = \frac{\xi}{2r^3}z' + \frac{\xi^2}{16r^4}z'', \tag{A.46}$$

$$\frac{1}{c}c'' - \frac{2}{c^2}c'c' - \frac{1}{r}\frac{1}{c}c' = \frac{\xi}{2r^3y}y' + \frac{\xi^2}{16r^4y}y'' - 2\frac{\xi}{4r^2y}y'\frac{\xi}{4r^2y}y' + \frac{\xi}{4r^3y}y',$$

and examine the particular case

$$c = \frac{1+\dfrac{\xi}{4r}}{1-\dfrac{\xi}{4r}}, \qquad \xi > 0. \tag{A.47}$$

We have

$$y' = 2\frac{1}{\left(1-\dfrac{\xi}{4r}\right)^2}, \qquad y'' = 4\frac{1}{\left(1-\dfrac{\xi}{4r}\right)^3}, \tag{A.48}$$

whence

$$\frac{1}{c}c'' - \frac{2}{c^2}c'c' - \frac{1}{r}\frac{1}{c}c' = 3\frac{\xi}{4r^3y}y' + \frac{\xi^2}{16r^4y}y'' - 2\frac{\xi}{4r^2y}y'\frac{\xi}{4r^2y}y'$$

$$= 6\frac{1}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)\left(1-\dfrac{\xi}{4r}\right)} + \frac{4}{r^2}\frac{\left(\dfrac{\xi}{4r}\right)^2}{\left(1+\dfrac{\xi}{4r}\right)\left(1-\dfrac{\xi}{4r}\right)^2} - \frac{8}{r^2}\frac{\left(\dfrac{\xi}{4r}\right)^2}{\left(1+\dfrac{\xi}{4r}\right)^2\left(1-\dfrac{\xi}{4r}\right)^2}$$

$$= 6\frac{1}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)\left(1-\dfrac{\xi}{4r}\right)} - \frac{4}{r^2}\frac{\left(\dfrac{\xi}{4r}\right)^2}{\left(1+\dfrac{\xi}{4r}\right)^2\left(1-\dfrac{\xi}{4r}\right)}$$

$$= -\frac{1}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)\left(1-\dfrac{\xi}{4r}\right)}\left[-6 + 4\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)}\right].$$

We shall confine the treatment to the relativistic $\epsilon = -1$:

$$m = -\left(1+\frac{\xi}{4r}\right)^4, \tag{A.49}$$

obtaining

$$\frac{m'}{m}+\frac{2}{r}=-4\frac{1}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)}+\frac{2}{r}$$

and

$$\frac{1}{2}\frac{c'}{c}\left(\frac{m'}{m}+\frac{2}{r}\right)=-\frac{1}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)\left(1-\frac{\xi}{4r}\right)}\left[-\frac{4}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)}+\frac{2}{r}\right]=-\frac{2}{r^2}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)^2},$$

$$\frac{m'}{m}\left(\frac{1}{4}\frac{m'}{m}+\frac{1}{r}\right)=-\frac{4}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)}\left[-\frac{1}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)}+\frac{1}{r}\right]=-\frac{4}{r^2}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)^2},$$

and also

$$\left(\frac{m'}{m}\right)'=-\frac{2}{r}\frac{m'}{m}+\frac{\xi}{4r^2}\frac{m'}{m}\frac{1}{\left(1+\frac{\xi}{4r}\right)}.$$

Therefore,

$$\frac{1}{m}\left(m''-\frac{1}{r}m'-\frac{3}{2}\frac{(m')^2}{m}\right)=\left(\frac{m'}{m}\right)'-\frac{1}{r}\frac{m'}{m}-\frac{1}{2}\frac{(m')^2}{m^2}$$

$$=\frac{m'}{m}\left[-\frac{2}{r}+\frac{\xi}{4r^2}\frac{1}{\left(1+\frac{\xi}{4r}\right)}-\frac{1}{r}+2\frac{\xi}{4r^2}\frac{1}{\left(1+\frac{\xi}{4r}\right)}\right]$$

$$=\frac{m'}{m}\frac{3}{r}\left[-1+\frac{\xi}{4r}\frac{1}{\left(1+\frac{\xi}{4r}\right)}\right]=-\frac{m'}{m}\frac{3}{r}\frac{1}{\left(1+\frac{\xi}{4r}\right)}=\frac{12}{r^2}\frac{\left(\frac{\xi}{4r}\right)}{\left(1+\frac{\xi}{4r}\right)^2},$$

whence we have

$$n_1=0. \tag{A.50}$$

Thereafter, we obtain

$$\frac{1}{c}c''-\frac{2}{c^2}c'c'-\frac{1}{r}\frac{1}{c}c'-\frac{c'}{c}\frac{m'}{m}=$$

$$-\frac{1}{r^2}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)\left(1-\frac{\xi}{4r}\right)}\left[-6+4\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)}\right]-\frac{2}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)\left(1-\frac{\xi}{4r}\right)}\frac{4}{r}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)}$$

$$= 6\frac{1}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)\left(1-\dfrac{\xi}{4r}\right)}\left[1-2\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)}\right] = 6\frac{1}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)^2},$$

whence

$$n_2 = n_3 = 0. \tag{A.51}$$

The conclusions (A.50) and (A.51) tell us that

$$a_n{}^i{}_{im} = 0 \quad \text{when} \quad N = 4. \tag{A.52}$$

Under these conditions, the tensor (A.39) reduces to

$$\begin{aligned}
a_n{}^i{}_{km} = \frac{2}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)^2}\Bigg[& 2\left(u_{mn}u_k{}^i - u_{kn}u_m{}^i\right) \\
& -\frac{3}{c^2}\left(n_n(n_kb_m - n_mb_k)b^i - \frac{1}{m}b_n(n_kb_m - n_mb_k)n^i\right) \\
& -3\Big(n_n(n_mu_k{}^i - n_ku_m{}^i) - (n_mu_{nk} - n_ku_{nm})n^i\Big) \\
& -\frac{1}{c^2}\left(\frac{1}{m}b_n(b_mu_k{}^i - b_ku_m{}^i) - (b_mu_{nk} - b_ku_{nm})b^i\right)\Bigg],
\end{aligned} \tag{A.53}$$

which can also be written as

$$\begin{aligned}
a_n{}^i{}_{km} = \frac{2}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)^2}\Bigg[& 2\left(u_{mn}u_k{}^i - u_{kn}u_m{}^i\right) \\
& -3\Big(n_n(n_mu_k{}^i - n_ku_m{}^i) - (n_mu_{nk} - n_ku_{nm})n^i\Big) \\
& -\frac{1}{c^2}\left(\frac{1}{m}b_n\Big(b_m(u_k{}^i - 3n_kn^i) - b_k(u_m{}^i - 3n_mn^i)\Big) - \Big(b_m(u_{nk} - 3n_nn_k) - b_k(u_{nm} - 3n_nn_m)\Big)b^i\right)\Bigg].
\end{aligned} \tag{A.54}$$

Contracting the last tensor by the vector b yields the simple result, namely,

$$b^m a_n{}^i{}_{km} = \frac{2}{r^2}\frac{\dfrac{\xi}{4r}}{\left(1+\dfrac{\xi}{4r}\right)^2}\left[-\left(\frac{1}{m}b_n(u_k{}^i - 3n_kn^i) - (u_{nk} - 3n_nn_k)b^i\right)\right] \tag{A.55}$$

and

$$b^n a_n{}^i{}_{km} = \frac{2}{r^2}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)^2}\left[-\frac{1}{m}\Big(b_m(u_k{}^i - 3n_k n^i) - b_k(u_m{}^i - 3n_m n^i)\Big)\right], \tag{A.56}$$

from which it follows that

$$(bb^m b^n - b^m y^n - b^n y^m) a_n{}^i{}_{km} =$$

$$\frac{2}{r^2}\frac{\frac{\xi}{4r}}{\left(1+\frac{\xi}{4r}\right)^2}\left[-(u_{nk} - 3n_n n_k) b^i y^n + \frac{1}{m} b(u_k{}^i - 3n_k n^i) - \frac{1}{m} b_k(u_m{}^i - 3n_m n^i) y^m\right]. \tag{A.57}$$

Appendix B: Curvature tensor of the space $\mathcal{FR}^{PD}_{g;c}$

Below the evaluations are restricted by the particular conditions

$$\nabla_n b_m - \nabla_m b_n = 0, \quad g = const, \tag{B.1}$$

simultaneously keeping the general case

$$c = c(x) \tag{B.2}$$

of the norm value $||b||$.

In terms of the convenient notation

$$(ys) = y^j y^h \nabla_j b_h \tag{B.3}$$

the spray coefficients of the space $\mathcal{FR}^{PD}_{g;c}$ read

$$G^i = \frac{g}{\nu}(ys) v^i + a^i{}_{km} y^m y^k \tag{B.4}$$

(see [1]), entailing

$$G^i{}_k = -\frac{1}{\nu}\nu_k \frac{g}{\nu}(ys) v^i + 2\frac{g}{\nu} s_k v^i + \frac{g}{\nu}(ys) r^i{}_k + 2a^i{}_{km} y^m, \tag{B.5}$$

where $s_k = y^h \nabla_k b_h$ and

$$\nu_k = \frac{1}{q} v_k + (1-c^2) g b_k. \tag{B.6}$$

The derivative

$$\frac{\partial\left(\frac{1}{\nu}\nu_k\right)}{\partial y^m} = -\frac{1}{\nu^2}\nu_k \nu_m + \frac{1}{\nu}\frac{1}{q}\eta_{km} \tag{B.7}$$

will be applied.

Differentiating (B.4) with respect to y^m yields the result

$$G^i{}_{km} = -\frac{1}{\nu}\nu_k\left(-\frac{1}{\nu}\nu_m\frac{g}{\nu}(ys)v^i + 2\frac{g}{\nu}s_m v^i + \frac{g}{\nu}(ys)r^i{}_m\right) - \left(-\frac{1}{\nu^2}\nu_k\nu_m + \frac{1}{\nu}\frac{1}{q}\eta_{km}\right)\frac{g}{\nu}(ys)v^i$$

$$-2\frac{g}{\nu^2}\nu_m s_k v^i + 2\frac{g}{\nu}(\nabla_m b_k)v^i + 2\frac{g}{\nu}s_k r^i{}_m$$

$$-\frac{g}{\nu^2}(ys)\nu_m r^i{}_k + 2\frac{g}{\nu}s_m r^i{}_k + \Delta,$$

which can conveniently be written as follows:

$$G^i{}_{km} = -\left(-\frac{2}{\nu^2}\nu_k\nu_m + \frac{1}{\nu}\frac{1}{q}\eta_{km}\right)\frac{g}{\nu}(ys)v^i + 2\frac{g}{\nu}(\nabla_m b_k)v^i$$

$$-2\frac{g}{\nu^2}(\nu_m s_k + \nu_k s_m)v^i + 2\frac{g}{\nu}(s_k r^i{}_m + s_m r^i{}_k) - \frac{g}{\nu^2}(ys)(\nu_m r^i{}_k + \nu_k r^i{}_m) + 2a^i{}_{km}. \tag{B.8}$$

Next, we perform required differentiation with respect to x^k, obtaining

$$2\frac{\partial \bar{G}^i}{\partial x^k} = \frac{\partial G^i}{\partial x^k} = -\frac{g}{\nu^2}(ys)\Big(q_{,k} + g(1-c^2)b_{j,k}y^j - 2gbcc_k\Big)v^i$$

$$+\frac{g}{\nu}v^i y^m y^n \nabla_k\nabla_m b_n - \frac{g}{\nu}(ys)(b_{j,k}y^j b^i + bb^i_{,k}) + \Delta.$$

Since

$$q_{,k} = -\frac{1}{q}bb_{j,k}y^j = -\frac{1}{q}bs_k + \Delta, \tag{B.9}$$

we can write

$$\frac{\partial G^i}{\partial x^k} = \frac{g}{q\nu^2}(ys)\Big(b - gq(1-c^2)\Big)s_k v^i + \frac{g}{\nu^2}(ys)2gbcc_k v^i$$

$$+\frac{g}{\nu}v^i y^m y^n \nabla_k\nabla_m b_n - \frac{g}{\nu}(ys)(s_k b^i + b\nabla_k b^i) + \Delta. \tag{B.10}$$

Also,

$$y^j\frac{\partial G^i}{\partial x^j} = \frac{g}{q\nu^2}(ys)\Big(b - gq(1-c^2)\Big)(ys)v^i + \frac{g}{\nu^2}(ys)2gbc(yc)v^i$$

$$+\frac{g}{\nu}v^i y^j y^m y^n \nabla_j\nabla_m b_n - \frac{g}{\nu}(ys)((ys)b^i + bs^i) + \Delta$$

with the vector

$$e_k = \frac{b}{q^2}v_k - b_k, \tag{B.11}$$

which obeys the equality

$$\frac{\partial e_k}{\partial y^j} = \frac{b}{q^2}\eta_{kj} - \frac{1}{q^2}v_k e_j.$$

After that, we derive the equality

$$\nu^2 \frac{\partial\Big(y^j\dfrac{\partial G^i}{\partial x^j}\Big)}{\partial y^k} = -2\frac{g}{q\nu}\nu_k(ys)\Big(b-gq(1-c^2)\Big)(ys)v^i - \frac{g}{q^3}v_k(ys)\Big(b-gq(1-c^2)\Big)(ys)v^i$$

$$+2\frac{g}{q}(ys)\Big(b-gq(1-c^2)\Big)s_kv^i + \frac{g}{q}(ys)\Big(b-gq(1-c^2)\Big)(ys){r^i}_k$$

$$+\frac{g}{q}(ys)\Big(b_k-g\frac{1}{q}v_k(1-c^2)\Big)(ys)v^i$$

$$-4\frac{1}{\nu}\nu_kg(ys)gbc(yc)v^i + g(ys)2gb_kc(yc)v^i + g(ys)2gbcc_kv^i$$

$$+g(ys)2gbc(yc){r^i}_k + gs_k2gbc(yc)v^i$$

$$-g\nu_kv^iy^jy^my^n\nabla_j\nabla_mb_n + g\nu y^jy^my^n\nabla_j\nabla_mb_n{r^i}_k$$

$$+g\nu v^i\Big[{\delta^j}_ky^my^n + 2y^jy^m{\delta^n}_k\Big]\nabla_j\nabla_mb_n$$

$$+g\nu_k(ys)((ys)b^i+bs^i) - g\nu s_k((ys)b^i+bs^i) - g\nu(ys)(s_kb^i+b_ks^i) + \Delta.$$

With these observations, we find

$$\nu^2\left[3\frac{\partial G^i}{\partial x^k} - \frac{\partial\Big(y^j\dfrac{\partial G^i}{\partial x^j}\Big)}{\partial y^k}\right] =$$

$$3\frac{g}{q}(ys)\Big(b-gq(1-c^2)\Big)s_kv^i + 3g(ys)2gbcc_kv^i$$

$$+3g\nu v^iy^my^n\nabla_k\nabla_mb_n - 3g\nu(ys)s_kb^i - 3g\nu(ys)b\nabla_kb^i$$

$$+2\frac{g}{q\nu}\nu_k(ys)\Big(b-gq(1-c^2)\Big)(ys)v^i + \frac{g}{q^3}v_k(ys)\Big(b-gq(1-c^2)\Big)(ys)v^i$$

$$-2\frac{g}{q}(ys)\Big(b-gq(1-c^2)\Big)s_kv^i - \frac{g}{q}(ys)\Big(b-gq(1-c^2)\Big)(ys){r^i}_k$$

$$-\frac{g}{q}(ys)\Big(b_k-g\frac{1}{q}v_k(1-c^2)\Big)(ys)v^i$$

$$+4\frac{1}{\nu}\nu_k g(ys)gbc(yc)v^i - g(ys)2gb_kc(yc)v^i - g(ys)2gbcc_kv^i$$

$$-g(ys)2gbc(yc){r^i}_k - gs_k2gbc(yc)v^i$$

$$+g\nu_kv^iy^jy^my^n\nabla_j\nabla_mb_n - g\nu y^jy^my^n\nabla_j\nabla_mb_n{r^i}_k$$

$$-g\nu v^i\Big[{\delta^j}_ky^my^n + 2y^jy^m{\delta^n}_k\Big]\nabla_j\nabla_mb_n$$

$$-g\nu_k(ys)((ys)b^i + bs^i) + g\nu s_k((ys)b^i + bs^i) + g\nu(ys)(s_kb^i + b_ks^i) + \Delta$$

$$= 2\frac{g}{q}(ys)\Big(b - gq(1-c^2)\Big)s_kv^i + 4g(ys)gbcc_kv^i$$

$$+2g\nu v^iy^my^n\nabla_k\nabla_mb_n - g\nu(ys)s_kb^i - 3g\nu(ys)b\nabla_kb^i$$

$$+2\frac{g}{q\nu}\nu_k(ys)\Big(b - gq(1-c^2)\Big)(ys)v^i + \frac{g}{q^3}v_k(ys)\Big(b - gq(1-c^2)\Big)(ys)v^i$$

$$-\frac{g}{q}(ys)\Big(b - gq(1-c^2)\Big)(ys){r^i}_k$$

$$-\frac{g}{q}(ys)\Big(b_k - g\frac{1}{q}v_k(1-c^2)\Big)(ys)v^i$$

$$+\left(4\frac{1}{\nu}\nu_kg(ys)gbcv^i - 2g(ys)gb_kcv^i - 2g(ys)gbc{r^i}_k - 2gs_kgbcv^i\right)(yc)$$

$$-g\Big(y^jy^my^n\nabla_j\nabla_mb_n\Big)(\nu{r^i}_k - \nu_kv^i) - 2g\nu\Big(y^jy^m\nabla_j\nabla_mb_k\Big)v^i$$

$$-g\nu_k(ys)(ys)b^i + \Big[-gb\nu_k(ys) + g\nu s_kb + g\nu(ys)b_k\Big]s^i + \Delta.$$

Finally,

$$\nu^2\left[3\frac{\partial G^i}{\partial x^k} - \frac{\partial\Big(y^j\frac{\partial G^i}{\partial x^j}\Big)}{\partial y^k}\right] =$$

$$-g\Big(y^jy^my^n\nabla_j\nabla_mb_n\Big)(\nu{r^i}_k - \nu_kv^i) + 2g\nu y^jy^m\Big(\nabla_k\nabla_mb_j - \nabla_j\nabla_mb_k\Big)v^i$$

$$-3g\nu(ys)b\nabla_k b^i + g\Big[\nu s_k - \nu_k(ys) + \frac{1}{b}\nu(ys)b_k\Big]\Big(bs^i - (ys)b^i\Big) + \frac{1}{b}g\nu(ys)(ys)b_kb^i$$

$$+\left(4\frac{1}{\nu}\nu_k g(ys)gbcv^i - 2g(ys)gb_kcv^i - 2g(ys)gbcr^i{}_k - 2gs_kgbcv^i\right)(yc) + 4g(ys)gbcc_kv^i$$

$$+2\frac{g}{q}(ys)\Big(b - gq(1-c^2)\Big)s_kv^i$$

$$+2\frac{g}{q\nu}(ys)\Big(b - gq(1-c^2)\Big)(ys)\nu_kv^i$$

$$-\frac{g}{q}(ys)\Big(b - gq(1-c^2)\Big)(ys)\eta^i{}_k$$

$$-\frac{g}{q}(ys)\Big(b_k - g\frac{1}{q}v_k(1-c^2)\Big)(ys)v^i + \Delta. \tag{B.12}$$

After that, we evaluate

$$\nu^4G^i{}_jG^j{}_k = \Big[-\nu_jg(ys)v^i + 2g\nu s_jv^i + g\nu(ys)r^i{}_j\Big]\Big[-\nu_kg(ys)v^j + 2g\nu s_kv^j + g\nu(ys)r^j{}_k\Big] =$$

$$-\nu_jg(ys)v^i\Big[-\nu_kg(ys)v^j + 2g\nu s_kv^j + g\nu(ys)r^j{}_k\Big]$$

$$+2g\nu s_jv^i\Big[-\nu_kg(ys)v^j + 2g\nu s_kv^j + g\nu(ys)r^j{}_k\Big]$$

$$+g\nu(ys)r^i{}_j\Big[-\nu_kg(ys)v^j + 2g\nu s_kv^j + g\nu(ys)r^j{}_k\Big]$$

and arrive at

$$2\nu^4G^mG^i{}_{km} = 2g(ys)v^m\left(2\nu_k\nu_m - \nu\frac{1}{q}\eta_{km}\right)g(ys)v^i + 2g(ys)v^m2g\nu^2(\nabla_mb_k)v^i$$

$$-2g\nu(ys)v^m2g(\nu_ms_k + \nu_ks_m)v^i + 2g\nu^2(ys)v^m2g(s_kr^i{}_m + s_mr^i{}_k)$$

$$-g\nu(ys)v^mg(ys)(\nu_mr^i{}_k + \nu_kr^i{}_m) + \Delta. \tag{B.13}$$

Therefore,

$$2\nu^4G^jG^i{}_{kj} - \nu^4G^i{}_jG^j{}_k =$$

$$2g^2(ys)v^m\left(2\nu_k\nu_m-\nu\frac{1}{q}\eta_{km}\right)(ys)v^i+4g^2(ys)v^m\nu^2(\nabla_m b_k)v^i$$

$$-4g^2\nu(ys)v^m(\nu_m s_k+\nu_k s_m)v^i+4g^2\nu^2(ys)v^m s_k r^i{}_m+4g^2\nu^2(ys)v^m s_m r^i{}_k$$

$$-g^2\nu(ys)v^m(ys)(\nu_m r^i{}_k+\nu_k r^i{}_m)$$

$$+\nu_j g^2(ys)\Big[-\nu_k(ys)v^j+2\nu s_k v^j+\nu(ys)r^j{}_k\Big]v^i$$

$$+2g^2\nu s_j\Big[\nu_k(ys)v^j-2\nu s_k v^j-\nu(ys)r^j{}_k\Big]v^i$$

$$+g^2\nu(ys)r^i{}_j\Big[\nu_k(ys)v^j-2\nu s_k v^j-\nu(ys)r^j{}_k\Big]+\Delta,$$

or, on reducing similar terms,

$$2\nu^4 G^j G^i{}_{kj}-\nu^4 G^i{}_j G^j{}_k=$$

$$2g^2(ys)v^m\left(2\nu_k\nu_m-\nu\frac{1}{q}\eta_{km}\right)(ys)v^i+4g^2(ys)v^m\nu^2(\nabla_m b_k)v^i$$

$$-2g^2\nu(ys)v^m(\nu_m s_k+\nu_k s_m)v^i+2g^2\nu^2(ys)v^m s_k r^i{}_m+4g^2\nu^2(ys)v^m s_m r^i{}_k$$

$$-g^2\nu(ys)v^m(ys)\nu_m r^i{}_k$$

$$+\nu_j g^2(ys)\Big[-\nu_k(ys)v^j+\nu(ys)r^j{}_k\Big]v^i$$

$$+2g^2\nu s_j\Big[-2\nu s_k v^j-\nu(ys)r^j{}_k\Big]v^i$$

$$-g^2\nu(ys)r^i{}_j\nu(ys)r^j{}_k+\Delta. \tag{B.14}$$

On applying the contractions

$$v^m s_m=(ys)-b\sigma, \qquad v^j r^i{}_j=v^i-(1-c^2)bb^i, \tag{B.15}$$

$$r^j{}_m r^i{}_j = r^i{}_m - (1-c^2)b^i b_m, \qquad v_j v^j = q^2 - (1-c^2)b^2, \tag{B.16}$$

and

$$\nu_j v^j = \nu - (1-c^2)\frac{1}{q}(b^2 + gc^2bq) \tag{B.17}$$

(see [1]), the above expression reads

$$\begin{aligned}
2\nu^4 G^j G^i{}_{kj} - \nu^4 G^i{}_j G^j{}_k = {} & 4g^2(ys)\nu_k\Big[\nu - (1-c^2)\frac{1}{q}(b^2 + gc^2bq)\Big](ys)v^i \\
& -2g^2(ys)\nu\frac{1}{q}\Big[v_k - (1-c^2)bb_k - \frac{1}{q^2}v_k(q^2 - (1-c^2)b^2)\Big](ys)v^i \\
& +4g^2(ys)\nu^2(s_k - b\sigma_k)v^i \\
& -2g^2\nu(ys)\Bigg(\Big[\nu - (1-c^2)\frac{1}{q}(b^2 + gc^2bq)\Big]s_k + \nu_k\Big((ys) - b\sigma\Big)\Bigg)v^i \\
& +2g^2\nu^2(ys)s_k\big[v^i - (1-c^2)bb^i\big] \\
& +4g^2\nu^2(ys)\big((ys) - b\sigma\big)r^i{}_k \\
& -g^2\nu(ys)(ys)\Big[\nu - (1-c^2)\frac{1}{q}(b^2 + gc^2bq)\Big]r^i{}_k \\
& +\nu_j g^2(ys)(ys)(\nu r^j{}_k - \nu_k v^j)v^i - 2g^2\nu s_j\Big[(ys)r^j{}_k + 2s_k v^j\Big]v^i \\
& - g^2\nu^2(ys)(ys)\Big[r^i{}_k - (1-c^2)b^i b_k\Big] + \Delta.
\end{aligned} \tag{B.18}$$

For the hh-curvature tensor $R^i{}_k$ we may use the formula

$$\begin{aligned}
K^2 R^i{}_k \ := {} & 2\frac{\partial \bar{G}^i}{\partial x^k} - \frac{\partial \bar{G}^i}{\partial y^j}\frac{\partial \bar{G}^j}{\partial y^k} - y^j\frac{\partial^2 \bar{G}^i}{\partial x^j \partial y^k} + 2\bar{G}^j\frac{\partial^2 \bar{G}^i}{\partial y^k \partial y^j} \\
= {} & 2\frac{\partial \bar{G}^i}{\partial x^k} - \bar{G}^i{}_j \bar{G}^j{}_k - y^j\frac{\partial \bar{G}^i{}_k}{\partial x^j} + 2\bar{G}^j \bar{G}^i{}_{kj}
\end{aligned} \tag{B.19}$$

(which is tantamount to the definition (3.8.7) on p. 66 of the book [7]); we apply the notation

$$\bar{G}^i = \frac{1}{2}G^i, \qquad \bar{G}^i{}_k = \frac{1}{2}G^i{}_k, \qquad \bar{G}^i{}_{km} = \frac{1}{2}G^i{}_{km}, \qquad \bar{G}^i{}_{kmn} = \frac{1}{2}G^i{}_{kmn}.$$

REFERENCES

[1] G. S. Asanov: Finsleroid-regular space developed. Berwald case, *arXiv:math.DG*/0711.4180v1 (2007).

[2] G. S. Asanov: Finsleroid–Finsler space with Berwald and Landsberg conditions, *arXiv:math.DG*/0603472 (2006).

[3] G. S. Asanov: Finsleroid–Finsler space and spray coefficients, *arXiv:math.DG*/0604526 (2006).

[4] G. S. Asanov: Finsleroid–Finsler spaces of positive–definite and relativistic types. *Rep. Math. Phys.* **58** (2006), 275–300.

[5] G. S. Asanov: Finsleroid–Finsler space and geodesic spray coefficients, *Publ. Math. Debrecen* **71/3-4** (2007), 397-412.

[6] G.S. Asanov: Finsleroid corrects pressure and energy of universe. Respective cosmological equations, *arXiv*:0707.3305 [math-ph] (2007).

[7] D. Bao, S. S. Chern, and Z. Shen: *An Introduction to Riemann-Finsler Geometry,* Springer, N.Y., Berlin (2000).

[8] J.L. Synge: *Relativity: the General Theory,* North–Holland, Amsterdam (1960).

[9] S. Weinberg: *Gravitation and Cosmology,* John Wiley, New York (1972).